

HATS-2b: A transiting extrasolar planet orbiting a K-type star showing starspot activity

M. Mohler-Fischer¹, L. Mancini¹, J. D. Hartman^{4,5}, G. Á. Bakos^{4,5,*,**}, K. Penev^{4,5}, D. Bayliss⁶, A. Jordán⁷, Z. Csubry^{4,5}, G. Zhou⁶, M. Rabus⁷, N. Nikolov², R. Brahm⁷, N. Espinoza⁷, L. A. Buchhave^{8,9}, B. Béky⁵, V. Suc⁷, B. Csák¹, T. Henning¹, D. J. Wright³, C. G. Tinney³, B. C. Addison³, B. Schmidt⁶, R. W. Noyes⁵, I. Papp¹⁰, J. Lázár¹⁰, P. Sári¹⁰, and P. Conroy⁶

¹ Max Planck Institute for Astronomy, Königstuhl 17, D-69117 Heidelberg, Germany
e-mail: mohler@mpia.de

² Astrophysics Group, School of Physics, University of Exeter, Stocker Road, EX4 4QL, Exeter, UK

³ Exoplanetary Science at UNSW, School of Physics, University of New South Wales, 2052, Australia and Australian Centre for Astrobiology, University of New South Wales, 2052, Australia

⁴ Department of Astrophysical Sciences, Princeton University, NJ 08544, USA

⁵ Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA

⁶ Research School of Astronomy and Astrophysics, Australian National University, Canberra, ACT 2611, Australia

⁷ Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, 7820436 Macul, Santiago, Chile

⁸ Niels Bohr Institute, University of Copenhagen, DK-2100, Copenhagen, Denmark

⁹ Centre for Star and Planet Formation, Natural History Museum of Denmark, University of Copenhagen, DK-1350 Copenhagen, Denmark

¹⁰ Hungarian Astronomical Association, Budapest, Hungary

Received ; Accepted

ABSTRACT

We report the discovery of HATS-2b, the second transiting extrasolar planet detected by the HATSouth survey. HATS-2b is moving on a circular orbit around a $V = 13.6$ mag, K-type dwarf star (GSC 6665-00236), at a separation of 0.0230 ± 0.0003 AU and with a period of 1.3541 days. The planetary parameters have been robustly determined using a simultaneous fit of the HATSouth, MPG/ESO 2.2 m/GROND, Faulkes Telescope South/Spectral transit photometry and MPG/ESO 2.2 m/FEROS, Euler 1.2 m/CORALIE, AAT 3.9 m/CYCLOPS radial-velocity measurements. HATS-2b has a mass of $1.37 \pm 0.16 M_J$, a radius of $1.14 \pm 0.03 R_J$ and an equilibrium temperature of 1567 ± 30 K. The host star has a mass of $0.88 \pm 0.04 M_\odot$, radius of $0.89 \pm 0.02 R_\odot$ and shows starspot activity. We characterized the stellar activity by analysing two photometric-follow-up transit light curves taken with the GROND instrument, both obtained simultaneously in four optical bands (covering the wavelength range of 3860 – 9520 Å). The two light curves contain anomalies compatible with starspots on the photosphere of the parent star along the same transit chord.

Key words. stars: planetary systems – stars: individual: (HATS-2, GSC 6665-00236) – stars: fundamental parameters – techniques: spectroscopic, photometric

1. Introduction

The first detection of a planet orbiting a main-sequence star (51 Peg; Mayor & Queloz 1995) started a new era of astronomy and planetary sciences. In the years since, the focus on exoplanetary discovery has steadily increased, resulting in more than 850 planets being detected in 677 planetary systems¹. Statistical implications of the exoplanet discoveries, based on different detection methods, have also been presented (e.g. Mayor et al. 2011; Howard et al. 2012; Cassan et al. 2012; Fressin et al. 2013). Most of these planets have been detected by the transit and radial velocity (RV) techniques. The former detects the decrease in a parent-star's brightness due to the transit of a planet in front of it, while the latter measures the Doppler shift of parent-star light

due to stellar motion around the star-planet barycenter. In the case of transiting extrasolar planets, the powerful combination of both methods permits a direct estimate of mass and radius of the planetary companion and therefore of the planetary average density and surface gravity. Such information is of fundamental importance in establishing the correct theoretical framework of planet formation and evolution (e.g. Liu et al. 2011; Mordasini et al. 2012a,b).

Thanks to the effectiveness of ground- and space-based transit surveys like TrES (Alonso et al. 2004), XO (McCullough et al. 2005), HATNet (e.g. Bakos et al. 2012a; Hartman et al. 2012), HATSouth (Penev et al. 2013), WASP (e.g. Hellier et al. 2012; Smalley et al. 2012), QES (Alsubai et al. 2011; Bryan et al. 2012), KELT (Siverd et al. 2012), COROT (e.g. Rouan et al. 2012; Pätzold et al. 2013) and Kepler (Borucki et al. 2011a,b; Batalha et al. 2012), one third of the transiting exoplanets known today were detected in the past 2 years. In some cases, exten-

* Alfred P. Sloan Research Fellow

** Packard Fellow

¹ exoplanet.eu, as at 2013, March 28

sive follow-up campaigns have been necessary to determine the correct physical properties of several planetary systems (e.g. Southworth et al. 2011; Barros et al. 2011; Mancini et al. 2013a), or have been used to discover other planets by measuring transit time variations (e.g. Steffen et al. 2013). With high-quality photometric observations it is also possible to detect transit anomalies which are connected with physical phenomena, such as star spots (Pont et al. 2007; Rabus et al. 2009; Désert 2011; Tregloan-Reed et al. 2013), gravity darkening (Barnes 2009; Szabó et al. 2011), stellar pulsations (Collier Cameron et al. 2010), tidal distortion (Li et al. 2010; Leconte et al. 2011), and the presence of additional bodies (exomoons) (Kipping et al. 2009; Tusnski & Valio 2011).

In this paper we report the detection of HATS-2b, the second confirmed exoplanet found by the HATSouth transit survey. HATSouth is the first global network of robotic wide-field telescopes, located at three sites in the Southern hemisphere: Las Campanas Observatory (Chile), Siding Spring Observatory (Australia) and H.E.S.S. (High Energy Stereoscopic System) site (Namibia). We refer the reader to Bakos et al. (2013), where the HATSouth instruments and operations are described in detail. HATS-2b is orbiting a K-type dwarf star and has characteristics similar to those of most hot-Jupiter detected so far. Photometric follow-up performed during two transits of this planet clearly reveals anomalies in the corresponding light curves, which are very likely related to the starspot activity of the parent star.

2. Observations

2.1. Photometry

The star HATS-2 (GSC 6665-00236; $V = 13.562 \pm 0.016$; J2000 $\alpha = 11^{\text{h}}46^{\text{m}}57^{\text{s}}.38$, $\delta = -22^{\circ}21'46''.77$, proper motion $\mu_{\alpha} = -45.8 \pm 1.1 \text{ mas/yr}$, $\mu_{\delta} = -1.3 \pm 1.5 \text{ mas/yr}$; UCAC4 catalogue, Zacharias et al. 2012) was identified as a potential exoplanet host based on photometry from all the instruments of the HATSouth facility (HS-1 to HS-6) between Jan 19 and Aug 10, 2010 (details are reported in Table 1). The detection light curve is shown in Figure 1. This figure shows that the discovery data is of sufficient quality that it permits fitting a standard limb darkened planetary transit model. A detailed overview of the observations, the data reduction and analysis is given in Bakos et al. (2013).

HATS-2 was afterwards photometrically followed-up three times by two different instruments at two different telescopes. On UT 2011, June 25, the mid-transit and the egress were observed with the “Spectral” imaging camera, mounted at the 2.0 m Faulkes Telescope South (FTS), at the Siding Spring Observatory (SSO). This partial transit permitted the refinement of the transit depth and ephemeris. The latter was particularly important for the subsequent follow-up observations performed with the MPG²/ESO 2.2m telescope at the La Silla Observatory (LSO). Two full transits were covered on February 28 and June 1, 2012, using GROND (Gamma-Ray Burst Optical/Near-Infrared Detector), which is an imaging camera capable of simultaneous photometric observations in four optical (identical to Sloan g , r , i , z) passbands (Greiner et al. 2008). The main characteristics of the cameras and details of the data reduction are described in Penev




Fig. 1. *Top panel:* Unbinned instrumental r -band light curve of HATS-2 folded with the period $P = 1.354133$ days resulting from the global fit described in Section 3. *Lower panel:* zoom in on the transit; the dark filled points show the light curve binned in phase using a bin size of 0.002. In both panels, the solid line shows the best-fit transit model (see Section 3.3).

et al. (2013). The GROND observations were performed with the telescope defocussed and using relatively long exposure times (80-90 s, 150-200 s cadence). This way minimises noise sources (e.g. flat-fielding errors, atmospheric variation or scintillation, variation in seeing, bad tracking and Poisson noise) and delivers high-precision photometry of transit events (Southworth et al. 2009). The light curves and their best-fitting models are shown in Fig. 2. Distortions in the GROND light curves are clearly visible, which we ascribe to stellar activity. These patterns are analysed in detail in Section 4. Table 1 gives an overview of all the photometric observations for HATS-2.


2.2. Spectroscopy

HATS-2 was spectroscopically followed-up between May 2011 and April 2012 by five different instruments at five individual telescopes. The follow-up observations started in May 2011 with high signal to noise (S/N) medium resolution ($\lambda/\Delta\lambda = 7000$) reconnaissance observations performed at the ANU 2.3 m telescope located at SSO, with the image slicing integral field spectrograph WiFeS (Dopita et al. 2007). The results showed no RV variation with amplitude greater than 2 km s^{-1} ; this excludes most false-positive scenarios involving eclipsing binaries. Furthermore, an initial determination of the stellar atmospheric parameters was possible ($T_{\text{eff},\star} = 4800 \pm 300 \text{ K}$, $\log g_{\star} = 4.4 \pm 0.3$), indicating that HATS-2 is a dwarf star. Within the same month, high precision RV follow-up observations started with the fibre-fed echelle spectrograph CORALIE (Queloz et al. 2000b) at the Swiss Leonard Euler 1.2 m telescope at LSO, followed by further high precision RV measurements obtained with the fibre-fed optical echelle spectrograph FEROS (Kaufer & Pasquini 1998) at the MPG/ESO 2.2 m telescope at LSO. Using the spectral synthesis code *SME* (‘Spectroscopy Made Easy’, Valenti & Piskunov 1996) on the FEROS spectra, it was possible to determine more accurate values for the stellar parameters (see

² Max Planck Gesellschaft

Table 1. Summary of photometric observations of HATS-2

Facility	Date(s)	# of images	Cadence (s)	Filter
Discovery				
HS-1 (Chile)	2010, Jan 24 - Aug 09	5913	280	Sloan <i>r</i>
HS-2 (Chile)	2010, Feb 11 - Aug 10	10195	280	Sloan <i>r</i>
HS-3 (Namibia)	2010, Feb 12 - Aug 10	1159	280	Sloan <i>r</i>
HS-4 (Namibia)	2010, Jan 26 - Aug 10	8405	280	Sloan <i>r</i>
HS-5 (Australia)	2010, Jan 19 - Aug 08	640	280	Sloan <i>r</i>
HS-6 (Australia)	2010, Aug 06	8	280	Sloan <i>r</i>
Follow-up				
FTS/Spectral	2011, June 25	158	60	Sloan <i>i</i>
MPG/ESO 2.2 m / GROND	2012, February 28	69	80	Sloan <i>g</i>
MPG/ESO 2.2 m / GROND	2012, February 28	70	80	Sloan <i>r</i>
MPG/ESO 2.2 m / GROND	2012, February 28	69	80	Sloan <i>i</i>
MPG/ESO 2.2 m / GROND	2012, February 28	71	80	Sloan <i>z</i>
MPG/ESO 2.2 m / GROND	2012, June 1	99	80	Sloan <i>g</i>
MPG/ESO 2.2 m / GROND	2012, June 1	99	80	Sloan <i>r</i>
MPG/ESO 2.2 m / GROND	2012, June 1	99	80	Sloan <i>i</i>
MPG/ESO 2.2 m / GROND	2012, June 1	99	80	Sloan <i>z</i>

**Fig. 2.** Phased transit light curves of HATS-2. The dates and instruments used for each event are indicated. The light curves are ordered according to the date and to the central wavelength of the filter used (Sloan *g*, *r*, *i* and *z*). Our best fit is shown by the solid lines (see Section 3.3). Residuals from the fits are displayed at the bottom, in the same order as the top curves. The GROND datapoints affected by anomalies are marked with red empty squares and were not considered in estimating the final values of the planetary-system parameters (see Sect. 3).

Sect. 3.1). Further RV measurements were obtained with the CYCLOPS fibre-based integral field unit, feeding the cross-dispersed echelle spectrograph UCLES, mounted at the 3.9 m Anglo-Australian Telescope (AAT) at SSO, and with the fibre-fed echelle spectrograph FIES at the 2.5 m telescope at the Nordic Optical Telescope in La Palma. We refer to Penev et al. (2013) for a more detailed description of the observations, the data reduction and the RV determination methods for each individual instrument that we utilized.

In total, 29 spectra were obtained, which are summarized in Table 2. Table 3 provides the high-precision RV and bisector span measurements. Figure 3 shows the combined high-precision RV measurements folded with the period of the transits. The error bars of the RV measurements include a component from astrophysical/instrumental jitter allowed to differ for the three instruments (Coralie: 74.0 ms^{-1} , FEROS: 44.0 ms^{-1} , CYCLOPS: 193.0 ms^{-1}).

3. Analysis

3.1. Stellar parameters

As already mentioned in Sect. 2.2, we estimated the stellar parameters, i.e. effective temperature T_{eff} , metallicity $[\text{Fe}/\text{H}]$, surface gravity $\log g$ and projected rotational velocity $v \sin i$, by applying *SME* on the high-resolution FEROS spectra. *SME* determines stellar and atomic parameters by fitting spectra from model atmospheres to observed spectra and estimates the parameter errors using the quality of the fit, expressed by the reduced χ^2 , as indicator. In case the S/N is not very high, or the spectrum is contaminated with telluric absorption features, cosmics or stellar emission lines, the reduced χ^2 does not always converge to unity, which leads to small errors for the stellar parameter values. To estimate of error bars, we used *SME* to determine the stellar parameters of each FEROS spectrum and calculated the weighted mean and corresponding scatter (weighted by the S/N of individual spectra). The results for the spectroscopic stellar parameters of the *SME* analysis are listed in Table 4.

By modeling the light curve alone it is possible to determine the stellar mean density, which is closely related

Table 2. Summary of spectroscopic observations of HATS-2. **Observing mode:** HPRV = high-precision RV measurements, RECON = reconnaissance observations.

Telescope/Instrument	Date Range	# of Observations	Instrument resolution	Observing mode
ANU 2.3 m/WiFeS	2011, May 10-15	5	7000	RECON
Euler 1.2 m/Coralie	2011, May 20-21	4	60000	HPRV
ESO/MPG 2.2 m/FEROS	2011, June 09-25	9	48000	HPRV
ESO/MPG 2.2 m/FEROS	2012, January 12	1	48000	HPRV
ESO/MPG 2.2 m/FEROS	2012, March 04-06	2	48000	HPRV
ESO/MPG 2.2 m/FEROS	2012, April 14-18	3	48000	HPRV
AAT 3.9 m/CYCLOPS	2012, January 05-12	4	70000	HPRV
NOT 2.5 m/FIES	2012, March 15	1	46000	RECON

Table 3. Relative RV and bisector span (BS) measurements of HATS-2 from various instruments used for high-precision RV measurements (c.f. Table 2). 5 data points determined with FEROS are not listed here and were not used for further analysis due to high error bars caused by bad weather conditions.

BJD (-2454000)	Relative RV (m s^{-1})	σ_{RV} (m s^{-1})	BS (m s^{-1})	σ_{BS} (m s^{-1})	Phase	Instrument
1701.52346	-90.37	33.00	-63.7	79.5	0.119	Coralie
1701.54622	-238.37	36.00	117.7	83.3	0.135	Coralie
1702.52760	168.63	33.00	31.3	58.6	0.860	Coralie
1702.55065	169.63	33.00	-30.8	74.2	0.877	Coralie
1721.50300	204.15	71.76	120.4	93.8	0.873	FEROS
1722.58300	371.99	100.77	-146.9	640.0	0.671	FEROS
1723.44500	-330.70	73.66	-211.0	207.2	0.307	FEROS
1736.46900	164.63	38.18	149.3	86.4	0.925	FEROS
1737.53800	261.69	119.02	-130.9	287.5	0.715	FEROS
1738.48600	-47.00	58.61	-222.1	146.9	0.415	FEROS
1932.22448	-33.63	21.80	1114.3	43.2	0.487	CYCLOPS
1933.21669	-40.53	51.20	3464.1	19.9	0.219	CYCLOPS
1934.12774	75.57	65.58	-4516.6	168.1	0.892	CYCLOPS
1938.81200	-337.82	139.13	-436.4	449.1	0.351	FEROS
1939.16016	47.07	49.44	9189.5	1723.7	0.608	CYCLOPS
1990.75600	294.99	103.71	46.3	201.6	0.711	FEROS
1992.82100	-335.90	72.26	-251.5	208.5	0.236	FEROS
2035.67400	31.55	77.56	270.6	223.1	0.882	FEROS

Notes. The Coralie RV uncertainties listed here are known to be underestimated. Updated estimates are available, but we list here the values we used in the analysis. We note that in any case a jitter is included in the analysis to account for any additional scatter to that implied by the uncertainties (cf. Table 5).

to the normalized semimajor axis a/R_* (Sect. 3.3) assuming a circular orbit. Furthermore, adding RV measurements allows the determination of these parameters for elliptical orbits as well.

To obtain the light curve model, quadratic limb-darkening coefficients are needed, which were determined using Claret (2004) and the initially determined stellar spectroscopic parameters. We used the Yonsei-Yale stellar evolution models (Yi et al. 2001; hereafter YY) to determine fundamental stellar parameters such as the mass, radius, age and luminosity. The light curve based stellar mean density and spectroscopy based effective temperature, coupled with isochrone analysis, together permit a more accurate stellar surface gravity determination. The newly determined value for $\log g_* = 4.50 \pm 0.05$ is consistent with the initial value of $\log g_* = 4.44 \pm 0.12$ thus we refrain from re-analyzing the spectra fixing the surface gravity to the revised value. The spectroscopic, photometric and derived stellar properties are listed in Table 4, whereas the adopted quadratic limb-darkening coefficients for the individual photometric filters are shown in Table 5.

To illustrate the position of HATS-2 in the H-R diagram, we plotted the normalized semi-major axis a/R_* ver-

sus effective temperature $T_{\text{eff},*}$. Figure 4 shows the values for HATS-2 with their 1- σ and 2- σ confidence ellipsoids as well as YY-isochrones calculated for the determined metallicity of $[\text{Fe}/\text{H}] = 0.15$ and interpolated to values between 1 and 14 Gyr in 1 Gyr increments from our adopted model.

3.2. Excluding blend scenarios

To rule out the possibility that HATS-2 is actually a blended stellar system mimicking a transiting planet system we conduct a detailed modeling of the light curves following the procedure described in Hartman et al. (2011). Based on this analysis we can reject hierarchical triple star systems with greater than 4.5σ confidence, and blends between a foreground star and a background eclipsing binary with $\sim 4\sigma$ confidence. Moreover, the only non-planetary blend scenarios which could plausibly fit the light curves (ones that cannot be rejected with greater than 5σ confidence) are scenarios which would have easily been rejected by the spectroscopic observations (these would be obviously double-lined systems, also yielding several km s^{-1} RV and/or BS variations). We thus conclude that the observed




Fig. 3. *Top panel:* high-precision RV measurements for HATS-2 from CORALIE (dark filled circles), FEROS (open triangles) and CYCLOPS (filled triangles) shown as a function of orbital phase, together with our best-fit model. Zero phase corresponds to the time of mid-transit. The center-of-mass velocity has been subtracted. *Second panel:* velocity O-C residuals from the best fit. The error bars include a component from astrophysical/instrumental jitter allowed to differ for the three instruments. *Third panel:* bisector spans (BS), with the mean value subtracted. Note the different vertical scales of the panels.




Fig. 4. YY-isochrones from Yi et al. (2001) for the metallicity of $[Fe/H] = 0.15$. Isochrones are plotted for ages between 1 and 14 Gyr in steps of 1 Gyr (left to right). The ellipses mark the 1- σ and 2- σ confidence ellipsoids for the determined values of a/R_* and T_{eff*} .

Table 4. Stellar parameters for HATS-2

Parameter	Value	Source
Spectroscopic properties		
T_{eff*} (K)	5227 ± 95	SME
$[Fe/H]$	0.15 ± 0.05	SME
$v \sin i$ (km s $^{-1}$)	1.5 ± 0.5	SME
$\log g_*$ (cgs)	4.44 ± 0.12	SME
Photometric properties		
V (mag)	13.562 ± 0.016	APASS ¹
B (mag)	14.490 ± 0.031	APASS
g (mag)	13.991 ± 0.012	APASS
r (mag)	13.260 ± 0.020	APASS
i (mag)	13.018 ± 0.021	APASS
J (mag)	11.906 ± 0.024	2MASS ²
H (mag)	11.475 ± 0.023	2MASS
K (mag)	11.386 ± 0.023	2MASS
Derived properties		
M_* ($M_\odot$)	0.882 ± 0.037	YY+a/ R_* +SME
R_* ($R_\odot$)	0.898 ± 0.019	YY+a/ R_* +SME
$\log g_*$ (cgs)	4.48 ± 0.02	YY+a/ R_* +SME
L_* ($L_\odot$)	0.54 ± 0.06	YY+a/ R_* +SME
M_V (mag)	5.61 ± 0.13	YY+a/ R_* +SME
M_K (mag)	3.64 ± 0.07	YY+a/ R_* +SME
Age (Gyr)	9.7 ± 2.9	YY+a/ R_* +SME
Distance (pc) ³	360 ± 11	YY+a/ R_* +SME

Notes. ¹ The AAVSO (A^American A^Asociation of V^Ariable S^Ars O^Bserver)s P^Hotometric A^Ll-S^Ky S^Urvey

² T^Wo M^Icron A^Ll S^Ky S^Urvey

³ A_V corrected

transit is caused by a planetary companion orbiting HATS-2.

3.3. Simultaneous analysis of photometry and radial velocity

Following the procedure described in Bakos et al. (2010), the FTS and GROND photometric follow-up measurements (Table 1) were simultaneously fitted with the high-precision RV measurements (Table 3). The light curve parameters, RV parameters, and planetary parameters are listed in Table 5.

Table 5 also contains values for the radial velocity jitter for all three instruments used for high-precision RV measurements. They are added in quadrature to the RV results of the particular instrument. These values are determined such that χ^2 per degree of freedom equals unity for each instrument when fitting a fiducial model. If χ^2 per degree of freedom is smaller than unity for that instrument, then no jitter would be added. The RV jitters are empirical numbers that are added to the measurements such that the actual scatter in the RV observations sets the posterior distributions on parameters like the RV semi-amplitude.

Allowing the orbital eccentricity to vary during the simultaneous fit, we include the uncertainty for this value in the other physical parameters. We find that the observations are consistent with a circular orbit ($e = 0.071 \pm 0.049$) and we therefore fix the eccentricity to zero for the rest of this analysis. Table 5 shows that the derived parameters obtained by including the distorted regions of the light curves are consistent with those derived with these regions excluded, indicating that the starspots themselves are not affecting the stellar or planet parameters in a significant way. We adopt the parameters obtained with the light curve distortions excluded in a fixed circular orbit.

Table 5. Orbital and planetary parameters for the HATS-2 system for different fitting scenarios: including the light curve distortions with free and fixed eccentricity e as well as excluding the light curve distortions with a fixed circular orbit. The last scenario was adopted for further analysis steps (parameters are highlighted in bold font).

Parameter	LC distortions included, $e \equiv 0$	LC distortions included, free e	LC distortions excluded, $e \equiv 0$
Light curve parameters			
P (days)	1.354133 ± 0.000001	1.354133 ± 0.000001	1.354133 ± 0.000001
T_c (BJD) ^a	$2455954.58576 \pm 0.00008$	$2455951.87748 \pm 0.00009$	$2455954.58576 \pm 0.00009$
T_{14} (days) ^a	0.0859 ± 0.0004	0.0859 ± 0.0004	0.0862 ± 0.0004
$T_{12} = T_{34}$ (days) ^a	0.0104 ± 0.0004	0.0107 ± 0.0004	0.0109 ± 0.0005
$a/R_\star$	$5.57^{+0.06}_{-0.09}$	5.65 ± 0.32	5.50 ± 0.09
$\zeta/R_\star$ ^b	26.48 ± 0.06	26.58 ± 0.07	26.52 ± 0.07
$R_p/R_\star$	0.1317 ± 0.0007	0.1326 ± 0.0008	0.1335 ± 0.0010
$b \equiv a \cos i/R_\star$	$0.214^{+0.061}_{-0.070}$	$0.265^{+0.053}_{-0.075}$	$0.271^{+0.055}_{-0.074}$
i (deg)	87.8 ± 0.7	87.4 ± 0.7	87.2 ± 0.7
Limb-darkening coefficients^c			
a_g (linear term)	0.7052	0.7052	0.7052
b_g (quadratic term)	0.1168	0.1168	0.1168
a_r	0.4756	0.4756	0.4756
b_r	0.2487	0.2487	0.2487
a_i	0.3617	0.3617	0.3617
b_i	0.2744	0.2744	0.2744
a_z	0.2861	0.2861	0.2861
b_z	0.2873	0.2873	0.2873
Radial velocity parameters			
K (ms ⁻¹)	272.2 ± 30.5	278.7 ± 33.0	268.9 ± 29.0
$e \cos \omega$	0.000	-0.033 ± 0.052	0.000
$e \sin \omega$	0.000	-0.023 ± 0.060	0.000
e	0.000	0.071 ± 0.049	0.000
ω (deg)	0.000	216 ± 77	0.000
RV jitter Coralie (ms ⁻¹)	74.0	74.0	74.0
RV jitter FEROS (ms ⁻¹)	44.0	44.0	44.0
RV jitter CYCLOPS (ms ⁻¹)	193.0	193.0	193.0
Planetary parameters			
M_p (M _J)	1.369 ± 0.158	1.397 ± 0.171	1.345 ± 0.150
R_p (R _J)	1.139 ± 0.025	1.131 ± 0.065	1.168 ± 0.030
$C(M_p, R_p)$ ^d	0.11	-0.26	0.08
ρ_p (g/cm ³)	1.15 ± 0.15	1.20 ± 0.28	1.05 ± 0.14
$\log g_p$ (cgs)	3.42 ± 0.05	3.43 ± 0.08	3.39 ± 0.05
a (AU)	0.0230 ± 0.0003	0.0230 ± 0.0003	0.0230 ± 0.0003
T_{eq} (K)	1567 ± 30	1554 ± 57	1577 ± 31
Θ^e	0.062 ± 0.007	0.064 ± 0.009	0.060 ± 0.007
$\langle F \rangle$ (10 ⁸ erg s ⁻¹ cm ⁻²) ^f	1.36 ± 0.11	1.32 ± 0.20	1.40 ± 0.11

Notes.

- ^a T_C : Reference epoch of mid transit that minimizes the correlation with the orbital period. BJD is calculated from UTC. T_{14} : total transit duration, time between first to last contact; $T_{12} = T_{34}$: ingress/egress time, time between first and second, or third and fourth, contact.
- ^b Reciprocal of the half duration of the transit used as a jump parameter in our MCMC analysis in place of $a/R_\star$. It is related to $a/R_\star$ by the expression $\zeta/R_\star = a/R_\star \cdot (2\pi(1 + e \sin \omega))/(P\sqrt{1 - b^2}\sqrt{1 - e^2})$ (Bakos et al. 2010).
- ^c Values for a quadratic law given separately for the Sloan g , r , i and z filters. These values were adopted from the tabulations by Claret (2004) according to the spectroscopic (SME) parameters listed in Table 4.
- ^d Correlation coefficient between the planetary mass M_p and radius R_p .
- ^e The Safronov number is given by $\Theta = (1/2)(V_{esc}/V_{orb})^2 = (a/R_p) \cdot (M_p/M_\star)$ (see Hansen & Barman 2007).
- ^f Incoming flux per unit surface area, averaged over the orbit.

4. Starspot analysis

Fig. 5 shows the combined four-colour GROND light curves for the two HATS-2 transit events that were observed with this imaging instrument. The slight difference in the transit depth among the datasets is due to the different wavelength range covered by each filter. In particular, the g , r , i and z filters are sensitive to wavelength ranges of 3860 – 5340 Å, 5380 – 7060 Å, 7160 – 8150 Å, and 8260 – 9520 Å, respectively.

4.1. Starspots and plages

From an inspection of Fig. 5, it is easy to note several distortions in the light curves. Such anomalies cannot be removed by choosing different comparison stars for the differential photometry, and we interpret them as the consequence of the planet crossing irregularities on the stellar photosphere, i.e. starspots. It is well known that starspots are at a lower temperature than the rest of the photosphere. The flux ratio should be therefore lower in the blue than the red. We




Fig. 5. Combined four-colour transit light curves of HATS-2 obtained with the GROND imaging system. Green dots are for the data taken in the g band, yellow upside down triangles for the r band, red squares for the i band, and purple triangles for the z band. *Top panel:* transit observed on February 28, 2012. The *bump* observed just after the midtransit is interpreted as the covering of a “cold” starspot by the planet. *Lower panel:* transit observed on June 1, 2012. In addition to the bump occurred near the egress part of the light curve, a “hot” spot manifested in the g band, just before the starting of the covering of the starspot.

thus expect to see stronger starspot features in the bluest bands.

The data taken on February 28, 2012, are plotted in the top panel of Fig. 5, where the *bump*, which is clearly present just after midtransit in all four optical bands, is explained by a starspot covered by the planet. In particular, considering the errorbars, the g , r , and i points in the starspot feature look as expected, whereas the feature in z is a bit peaked, especially the highest points at the peak of the bump at roughly BJD(TDB) 2455985.735. Before the starspot feature, it is also possible to note that the fluxes measured in the g and r bands are lower than those in the other two reddest bands, as if the planet were occulting a hotter zone of the stellar chromosphere. Actually, the most sensitive indicators of the chromospheric activity of a star in the visible spectrum are the emission lines of Ca II H&K3968, Kλ3933, and Hα λ6563, which in our case fall on the transmission wings of the g and r GROND passbands.

Within the transit observed on June 1, 2012, whose data are plotted in the lower panel of Fig. 5, we detected another starspot, which occurred near the transit-egress zone of the light curve. Again, before the starspot feature, we note another “hotspot” in the g band, which has its peak at roughly BJD(TDB) 2456079.681.

While these hotspot distortions could be caused by differential color extinction or other time-correlated errors (i.e. red noise) of atmospheric or astrophysical origin, there is also a physical interpretation that seems potentially plausible, i.e. that of a “plage”. A plage is a chromospheric region typically located near active starspots, and usually forming before the starspots appear, and disappearing after the starspots vanish from a particular area. Accordingly, a plage occurs most often near a starspot region. As a matter of fact, in the GROND light curves, our plages are located just before each starspot. One can argue that the plage in the second transit is visible only in the g band, but this

can be explained by temperature fluctuations in the chromosphere, which causes a lack of ionized hydrogen, and by the fact that the Ca II lines are much stronger than the H α line for a K-type star like HATS-2. Another argument supporting this *plage-starspot* scenario is that, for these old stars, a solar-like relation between photospheric and chromospheric cycles is expected, the photospheric brightness varying in phase with that of the chromosphere (Lockwood et al. 2007).

We note, however, that if these are plages they must be rather different from solar plages, which are essentially invisible in broad-band optical filters unless they are very close to the solar limb. Detecting a plage feature through a broad-band filter near the stellar center suggests a much larger temperature contrast and/or column density of chromospheric gas than in the solar case.

4.2. Modelling transits and starspots

We modelled the GROND transit light curves of HATS-2 with the PRISM³ and GEMC codes (Tregloan-Reed et al. 2013). The first code models a planetary transit over a spotted star, while the latter one is an optimisation algorithm for finding the global best fit and associated uncertainties. Using these codes, one can determine, besides the ratio of the radii $R_p/R_\star$, the sum of the fractional radii, $r_p + r_\star = (R_p + R_\star)/a$, the limb darkening coefficients, the transit midpoint T_0 , and the orbital inclination i , as well as the photometric parameters of the spots, i.e. the projected longitude and the latitude of their centres (θ and ϕ), the spot size r_{spot} and the spot contrast ρ_{spot} , which is basically the ratio of the surface brightness of the spot to that of the surrounding photosphere. Unfortunately, the current versions of PRISM and GEMC are set to fit only a single starspot (or hotspot), so we excluded the g -band dataset of the 2nd transit from the analysis, because it contains a hotspot with high contrast ratio between stellar photosphere and spot, which strongly interferes with the best-fitting model for the light curve.

Given that the codes do not allow the datasets to be fitted simultaneously, we proceeded as follows. First, we modelled the seven datasets (1st transit: g, r, i, z ; 2nd transit: r, i, z) of HATS-2 separately; this step allowed us to restrict the search space for each parameter. Then, we combined the four light curves of the first transit into a single dataset by taking the mean value at each point from the four bands at that point and we fitted the corresponding light curve; this second step was necessary to find a common value for T_0, i, θ and ϕ . Finally, we fitted each light curve separately fixing the starspot position, the midtime of transit T_0 and the system inclination to the values found in the previous combined fit.

The light curves and their best-fitting models are shown in Fig. 6, while the derived photometric parameters for each light curve are reported in Table 6, together with the results of the MCMC error analysis for each solution.

Comparing Table 5 with Table 6 we find that the fitted light curve parameters from the analysis described in Section 3.3 are consistent with the parameters that result from the GEMC+PRISM model, except for the inclination which differs by more than 2σ . As already discussed in

Section 3.3, the joint-fit analysis was performed both considering and without considering the points contaminated by the starspots, and the results are consistent with each other. So, our conclusion is that the spots themselves are not systematically affecting the stellar or planet parameters in a significant way; the differences in the inclination between GEMC and our joint fit are most likely due to differences in the modeling.

4.3. Starspots discussion

The final value for the starspots angular radii comes from the weighted mean of the results in each band and is $r_{\text{spot}} = 9.02^\circ \pm 0.30^\circ$ for the starspot in the 1st transit (spot #1) and $r_{\text{spot}} = 19.16^\circ \pm 1.08^\circ$ for the starspot in the 2nd transit (spot #2), with a reduced χ^2_ν of 0.78 and 0.49 respectively, indicating a good agreement between the various light curves in each of the two transits. We note that the error of the angular size of the spot #2 is greater than that of the spot #1. While it may be that spot #2 is larger, we caution that its position near the limb of the star makes its size poorly constrained.

The above numbers translate to radii of $13\,145 \pm 552$ km and $27\,923 \pm 1729$ km, which are equivalent 0.05% and 0.2% of the stellar disk, respectively. Starspot sizes are in general estimated by doppler-imaging reconstructions (i.e. Collier Cameron 1992; Vogt et al. 1999) and their range is 0.1% to 22% of a stellar hemisphere, the inferior value being the detection limit of this technique (Strassmeier 2009). Our measurements are thus perfectly reasonable for a common starspot or for a starspot *assembly*, and in agreement with what has been found in other K-type stars.

Starspots are also interesting in terms of how the contrast changes with passband. In particular, we expect that moving from ultraviolet (UV) to infrared (IR) wavelengths the spot becomes brighter relative to the photosphere. Considering the starspot #2, its contrast decreases from r to z , even though this variation is inside the 1σ error (see Table 6). Considering that HATS-2 has an effective temperature $T_{\text{eff}\star} = 5227 \pm 95$ K and modeling both the photosphere and the starspot as blackbodies (Rabus et al. 2009; Sanchis-Ojeda & Winn 2011), we used Eq. (1) of Silva (2003) to estimate the temperature of the starspot #2 in each band, obtaining the following values: $T_{\text{spot}\#2,r} = 4916 \pm 105$ K, $T_{\text{spot}\#2,i} = 4895 \pm 121$ K and $T_{\text{spot}\#2,z} = 4856 \pm 120$ K. The weighted mean is $T_{\text{spot}\#2} = 4891.5 \pm 66.2$ K.

Unlike starspot #2, the spot contrasts for starspot #1 are inconsistent with expectations. The spot is too bright in r relative to the other bandpasses, and too faint in z . If we estimate the starspot temperature in each band, we find $T_{\text{spot}\#1,g} = 4345 \pm 97$ K, $T_{\text{spot}\#1,r} = 4604 \pm 109$ K, $T_{\text{spot}\#1,i} = 4318 \pm 128$ K and $T_{\text{spot}\#1,z} = 3595 \pm 180$ K. While the temperature in r is in agreement with those of g and i at $1 - 2\sigma$ level, and slight differences could be explained by chromospheric contamination (filaments, spicules, etc.), the temperature in z seems physically inexplicable. This effect is essentially caused by the z points at the peak of the starspot, at phase ~ 0.004 (see Fig. 6), which are higher than the other points. However, one has also to consider that errorbars in this band are larger than those found in the other bands. This is due to the fact that, since the GROND system design does not permit

³ PRISM (Planetary Retrospective Integrated Star-spot Model).

Table 6. Photometric parameters derived from the GEMC fitting of the GROND transit light curves.

1 st transit		<i>g</i>	<i>r</i>	<i>i</i>	<i>z</i>
Parameter	Symbol				
Radius ratio	$R_p/R_\star$	0.1348 ± 0.0011	0.1324 ± 0.0011	0.13145 ± 0.00096	0.1352 ± 0.0010
Sum of fractional radii	$r_p + r_\star$	0.2204 ± 0.0018	0.2232 ± 0.0018	0.2149 ± 0.0016	0.2161 ± 0.0015
Linear LD coefficient	u_1	0.749 ± 0.060	0.593 ± 0.051	0.352 ± 0.057	0.298 ± 0.039
Quadratic LD coefficient	u_2	0.171 ± 0.018	0.296 ± 0.043	0.218 ± 0.028	0.144 ± 0.024
Inclination (degrees) ^a	i	85.26 ± 0.40	85.26 ± 0.40	85.26 ± 0.40	85.26 ± 0.40
Longitude of spot (degrees) ^{a,b}	θ	5.78 ± 0.45	5.78 ± 0.45	5.78 ± 0.45	5.78 ± 0.45
Latitude of Spot (degrees) ^{a,c}	ϕ	76.52 ± 1.94	76.52 ± 1.94	76.52 ± 1.94	76.52 ± 1.94
Spot angular radius (degrees) ^d	r_{spot}	8.85 ± 0.61	10.01 ± 0.72	8.93 ± 0.65	8.72 ± 0.50
Spot contrast ^e	ρ_{spot}	0.304 ± 0.035	0.546 ± 0.048	0.464 ± 0.052	0.251 ± 0.52
2 nd transit					
Radius ratio	$R_p/R_\star$	—	0.1356 ± 0.0012	0.13411 ± 0.00093	0.1307 ± 0.0011
Sum of fractional radii	$r_p + r_\star$	—	0.2108 ± 0.0019	0.2022 ± 0.0012	0.2039 ± 0.0017
Linear LD coefficient	u_1	—	0.473 ± 0.057	0.399 ± 0.049	0.252 ± 0.044
Quadratic LD coefficient	u_2	—	0.250 ± 0.038	0.230 ± 0.025	0.316 ± 0.049
Inclination (degrees) ^a	i	—	85.89 ± 0.40	85.89 ± 0.40	85.89 ± 0.40
Longitude of spot (degrees) ^{a,b}	θ	—	35.26 ± 1.20	35.26 ± 1.20	35.26 ± 1.20
Latitude of Spot (degrees) ^{a,c}	ϕ	—	80.60 ± 2.10	80.60 ± 2.10	80.60 ± 2.10
Spot angular radius (degrees) ^d	r_{spot}	—	20.14 ± 1.49	17.79 ± 2.17	18.28 ± 2.31
Spot contrast ^e	ρ_{spot}	—	0.753 ± 0.046	0.780 ± 0.054	0.789 ± 0.047

Notes.

^a This is a common value and was found from the preceding fit of the combined data (see text).

^b The longitude of the centre of the spot is defined to be 0° at the centre of the stellar disc and can vary from -90° to 90° .

^c The latitude of the centre of the spot is defined to be 0° at the north pole and 180° at the south pole.

^d Note that 90° degrees covers half of stellar surface.

^e Note that 1.0 equals to the surrounding photosphere.

to chose different exposure times for each band, we are forced to optimize the observations for the *r* and *i* bands. Consequently, considering both the filter-transmission efficiency and the color and the magnitude of HATS-2, the SNR in these two bands is better than that in *z*, for which we have larger uncertainty in the photometry. Taking these considerations into account, we estimated the final temperature of starspot #1 neglecting the *z*-band value, and obtaining $T_{\text{spot}\#1} = 4425 \pm 63$ K. In Fig. 7 the final values of the temperature contrast of the two starspots are compared with those of a sample of dwarf stars, which was reported by Berdyugina (2005). The derived contrast for the HATS-2 starspots is consistent with what is observed for other stars. As already observed by Strassmeier (2009), the temperature difference between photosphere and starspots can be not so different for stars of different spectral types. Moreover, in the case of long lifetime, the same starspot could be seen at quite different temperature (Kang & Wilson 1989). It is then very difficult to find any clear correlation between starspot temperatures and spectral classes of stars.

The achieved longitudes of the starspots are in agreement with a visual inspection of the light curves. The latitude of starspot #1, $76.52^\circ \pm 1.94^\circ$, matches well with that of starspot #2, $80.6^\circ \pm 2.1^\circ$, the difference being within 1σ .

Multiple planetary transits across the same spot complex can be used to constrain the alignment between the orbital axis of the planet and the spin axis of the star (e.g. Sanchis-Ojeda et al. (2011)). Unfortunately, from only two transits separated by 94 days we cannot tell whether or not the observed anomalies are due to the same complex. It is possible that they are. Following Solanki (2003) we esti-




Fig. 7. Spot temperature contrast with respect to the photospheric temperature in several dwarf stars. Gray square is from spot #2, black circle from spot #1, open circles from Berdyugina (2005). The name of the star and its spectral type is also reported for most of them. Note that some stars appear two times.

mate a typical lifetime of ~ 130 days for spots of the size seen here. Moreover the rotation period of $P_{\text{rot}} = 31 \pm 1$ d inferred assuming they are the same spot is consistent with the value of $P_{\text{rot}} = (30 \pm 10 \text{ d}) \sin i_\star$ estimated from the spectroscopically-determined sky-projected equatorial rotation speed. If they are the same spot complex, then the sky-projected spin-orbit alignment is $\lambda = 8^\circ \pm 8^\circ$, which is consistent with zero. We caution, however, that this value depends entirely on this assumption which could easily be




Fig. 6. Phased GROND light curves of HATS-2b compared to the best GEMC fits. The light curves and the residuals are ordered according to the central wavelength of the filter used. *Left panel:* transit observed on February 28, 2012. *Right panel:* transit observed on June 1, 2012; due to presence of the hotspot, the *g* band was not analysed with GEMC (see text).

wrong. Continued photometric monitoring of HATS-2, or spectroscopic observations of the Rossiter-McLaughlin effect, are necessary to measure the spin-orbit alignment of this system.

5. Conclusions

In this paper we have presented HATS-2b, the second planet discovered by the HATSouth survey. This survey is a global network of six identical telescopes located at three different sites in the Southern hemisphere (Bakos et al. 2013). The parameters of the planetary system were estimated by an accurate joint fit of follow-up RV and photometric measurements. In particular, we found that HATS-2b has a mass of $1.369 \pm 0.158 M_J$, and a radius of $1.139 \pm 0.025 R_J$. To set this target in the context of other transit planet detections, we plotted 4 different types of correlation diagrams for the population of transiting planets (Fig. 8). We analysed the location of determined parameters for HATS-2b and its host star HATS-2 in the following parameter spaces: planetary radius R_P vs. stellar radius R_* , planetary mass M_P vs. planetary equilibrium temper-

ature $T_{eq,P}$, planetary radius R_P vs. planetary equilibrium temperature $T_{eq,P}$, planetary radius R_P vs. stellar effective temperature $T_{eff,*}$, and planetary radius R_P vs. planetary mass M_P . As illustrated in Fig. 8, the analysed parameter relations lie well within the global distribution of known exoplanets.

Within each correlation diagram, at least one well characterized exoplanet can be found, whose parameters are consistent with those of HATS-2(b) within the error bars. Looking on the correlation between planetary and stellar radius, HATS-2(b) is almost like HAT-P-37(b). Comparing the planetary equilibrium temperature and planetary mass, HATS-2b is similar to TrES-2b. The relation between planetary equilibrium temperature and planetary mass shows an agreement with WASP-32b, while the relation between stellar effective temperature and planetary radius points out that HATS-2b agrees well with WASP-45b within the error bars. The focus on the planetary parameters radius and mass reveals a similarity to the transiting planet TrES-2. Comparing the atmospheres of exoplanets with similar physical parameters will be especially important to pursue




Fig. 8. Correlation diagrams for confirmed transit planets (exoplanet.eu, last updated Jan 10 2013). From top to bottom: planetary radius R_P vs. stellar radius R_* , planetary mass M_P vs. planetary equilibrium temperature $T_{eq,P}$, planetary radius R_P vs. planetary equilibrium temperature $T_{eq,P}$, planetary radius R_P vs. stellar effective temperature $T_{eff,*}$ and planetary radius R_P vs. planetary mass M_P . The position of the HATS-2 and HATS-2b parameters, respectively, are marked in red squares. Left panels give a global overview of the position of HATS-2 and HATS-2b in the population of known transit planets (black filled circles). Right panels illustrate a zoom in; green diamonds represent known exoplanets which fall within the errorbars of HATS-2(b) parameters. Short cuts: HP = HAT-P, W = WASP, T = TrES.

with e.g. the future ECHO space mission (Tinetti et al. 2012).

Very interesting is the detection of anomalies in the two multi-band photometric-follow-up light curves obtained with the GROND imaging instrument. We recognize the anomalies as starspots covered by HATS-2b during the two transit events, and used PRISM and GEMC codes (Tregloan-Reed et al. 2013) to re-fit the transit light curves, measuring the parameters of the spots. Both the starspots appear to have associated hot-spots, which appeared in the 1st transit in the *g* and *r* bands, and only in the *g* band in the 2nd transit. These hotspots could be physically interpreted as chromospheric active regions known as *plages*, which can be seen only in the GROND's bluest bands. We estimated the size and the temperature of the two starspots, finding values which are in agreement with those found in other G-K dwarf stars.

Acknowledgements. Development of the HATSouth project was funded by NSF MRI grant NSF/AST-0723074, operations are supported by NASA grant NNX09AB29G, and follow-up observations receive partial support from grant NSF/AST-1108686. Data presented in this paper is based partly on observations obtained with the HATSouth Station at the Las Campanas Observatory of the Carnegie Institution of Washington. This work is based on observations collected at the MPG/ESO 2.2m Telescope located at the ESO Observatory in La Silla (Chile), under programme IDs P087.A-9014(A), P088.A-9008(A), P089.A-9008(A), 089.A-9006(A) and Chilean time P087.C-0508(A). Operations of this telescope are jointly performed by the Max Planck Gesellschaft and the European Southern Observatory. GROND has been built by the high-energy group of MPE in collaboration with the LSW Tautenburg and ESO, and is operating as a PI-instrument at the MPG/ESO 2.2m telescope. We thank Timo Anguita and Régis Lachaume for their technical assistance during the observations at the MPG/ESO 2.2m Telescope. A.J. acknowledges support from Fondecyt project 1130857, Anillo ACT-086, BASAL CATA PFB-06 and the Millennium Science Initiative, Chilean Ministry of Economy (Nucleus P10-022-F). V.S. acknowledges support from BASAL CATA PFB-06. R.B. and N.E. acknowledge support from Fondecyt project 1095213. N.N. acknowledges support from an STFC consolidated grant. M.R. acknowledges support from FONDECYT postdoctoral fellowship N°3120097. L.M. thanks Jeremy Tregloan-Reed for his help in using of the PRISM and GEMC codes, and John Southworth and Valerio Bozza for useful discussions.

This paper also uses observations obtained with facilities of the Las Cumbres Observatory Global Telescope. Work at the Australian National University is supported by ARC Laureate Fellowship Grant FL0992131. We acknowledge the use of the AAVSO Photometric All-Sky Survey (APASS), funded by the Robert Martin Ayers Sciences Fund, and the SIMBAD database, operated at CDS, Strasbourg, France. Work at UNSW has been supported by ARC Australian Professorial Fellowship grant DP0774000, ARC LIEF grant LE0989347 and ARC Super Science Fellowships FS100100046.

References

- Alsubai, K. A., Parley, N. R., Bramich, D. M., et al. 2011, *MNRAS*, 417, 709
- Armitage, P. J. & Bonnell, I. A. 2002, *MNRAS*, 330, L11
- Bakos, G. Á., Torres, G., Pál, A., et al. 2010, *ApJ*, 710, 1724
- Bakos, G. Á., Hartman, J. D., Torres, G., et al. 2012, *AJ*, 144, 19
- Bakos, G. Á., Csabry, Z., Oenev, K., et al. 2013, *PASP*, 125, 0
- Ballester, G. E., Sing, D. K., & Herbert, F. 2007, *Nature*, 445, 511
- Barnes, J. W. 2009, *ApJ*, 705, 683
- Barros, S. C. C., Pollacco, D. L., Gibson, N. P., et al. 2004, *MNRAS*, 416, 2593
- Batalha, N. M., Rowe, J. F., Bryson, S. T., et al. 2012, *ApJS*, 204, 24
- Bean, J. L., Miller-Ricci Kempton, E. & Homeier, D. 2010, *Nature*, 468, 669
- Berdugina, S. V. 2005, *Living Rev. Solar Phys.*, 2, 8
- Bonomo, A. S., & Lanza, A. F. 2012, *A&A*, 547, A37
- Borucki, W. J., Koch, D., Jenkins, J., et al. 2009, *Science*, 325, 709
- Borucki, W. J., Koch, D. G., Basri, G., et al. 2011a, *ApJ*, 728, 117
- Borucki, W. J., Koch, D. G., Basri, G., et al. 2011b, *ApJ*, 736, 19
- Bryan, M. L., Alsubai, K. A., Latham, D. W., et al. 2012, *ApJ*, 750, 84
- Cassan, A., Kubas, D., Beaulieu, J.-P., et al. 2012, *Nature*, 481, 167
- Claret, A. 2004, *A&A*, 428, 1001
- Collier Cameron, A. 1992, *Surface inhomogeneities on late-type stars*, ed. P. B. Byrne, & D. J. Mullan (Springer, Berlin), 33
- Collier Cameron, A., Guenther, E., Smalley, B., McDonald, I. 2010, *MNRAS*, 407, 507
- D'Angelo, G., Durisen, R. H., Lissauer, J. J. 2011, *Exoplanets*, edited by S. Seager. (University of Arizona Press), p. 319
- Désert, J.-M., Charbonneau, D., Demory, B.-O., et al. 2011, *ApJS*, 197, 14
- Dopita, M., Hart, J., McGregor, P., et al. 2007, *Ap&SS*, 310, 255
- Fortney, J. J., Lodders, K., Marley, M. S., & Freedman, R. S. 2008, *ApJ*, 678, 1419
- Fressin, F., Torres, G., Charbonneau, D., et al. 2013, to appear in *ApJ*, arXiv:1301.0842
- Gaudi, B. S. & Winn, J. N. 2007, *ApJ*, 655, 550
- Greiner, J., Bornemann, W., Clemens, C., et al. 2008, *PASP*, 120, 405
- Gurdemir, L., Redfield, S., & Cuntz, M. 2012, *PASP*, 124, 141
- Hansen, B. M. S., & Barman, T. 2007, *ApJ*, 671, 861
- Hartman, J. D., Bakos, G. Á., Béky, B., et al. 2012, *AJ*, 144, 139
- Hartman, J. D., Bakos, G. Á., Torres, G., et al. 2011, *ApJ*, 742, 59
- Hellier, C., Anderson, D. R., Collier Cameron, A., et al. 2012, *MNRAS*, 426, 739
- Hirano, T., Narita, N., Sato, B. et al. 2012, *ApJ*, 759, L36
- Howard, A. W., Marcy, G. W., Bryson, S. T., et al. 2012, *ApJS*, 201, 15
- Hussain, G. A. J. 2002, *Astron. Nachr.*, 323, 349
- Kang, Y. W. & Wilson, R. E. 1989, *AJ*, 97, 848
- Kaufer, A., & Pasquini, L. 1998, *Proc. SPIE*, 3355, 844
- Kipping, D. M., Fossey, S. J., Campanella, G. 2009, *MNRAS*, 400, 398
- Knutson, H. A., Charbonneau, D., Allen, L. E., et al. 2007, *Nature*, 447, 183
- Leconte, J., Lai, D., Chabrier, G. 2011, *A&A*, 528, A41
- Li, S.-L., Miller, N. Lin, D. N. C., Fortney, J. J. 2010, *Nature*, 463, 1054
- Liu, H., Zhou, J.-L., Wang, S. 2011, *ApJ*, 732, 66
- Lockwood, G. W., Skiff, B. A., Henry, G. W., et al. 2007, *ApJS*, 171, 260
- Lubow, S. H., & Ida, S. 2011, *Exoplanets*, edited by S. Seager. (University of Arizona Press), p. 347
- Mancini, L., Southworth, J., Ciceri, S., et al. 2013, *A&A*, 551, A11
- Mancini, L., Nikolov, N., Southworth, J., et al. 2013, to appear in *MNRAS*, arXiv:1301.3005
- Mayor, M. & Queloz, D. 1995, *Nature*, 378, 355
- Mayor, M., Marmier, M., Lovi, C., et al. 2011, arXiv:1109.2497
- Mordasini, C., Alibert, Y., Klahr, H., Henning, T. 2012, *A&A*, 547, A111
- Mordasini, C., Alibert, Y., Georgy, C., et al. 2012, *A&A*, 547, A112
- Orosz, J. A., Welsh, W. F., Carter, J. A. et al. 2012a, *ApJ*, 758, 87
- Orosz, J. A., Welsh, W. F., Carter, J. A. et al. 2012b, *Science*, 337, 1511
- Pätzold, M., Endl, M., Csizmadia, Sz., et al. 2012, 545, A6
- Penev, K., Bakos, G. Á., Bayliss, D., et al. 2013, *AJ*, 145, 5
- Petrovay, K., & van Driel-Gesztelyi, L. 1997, *Sol. Phys.*, 176, 249
- Pont, F., Gilliland, R. L., Moutou, C., et al. 2007, *A&A*, 476, 1347
- Queloz, D., Eggenberger, A., Mayor, M., Perrier, C., Beuzit, J.-L., Naef, D., Sivan, J.-P., Udry, S. 2000, *A&A*, 359, L13
- Queloz, D., Mayor, M., Weber, L. et al. 2000, *A&A*, 354, 99
- Rabus, M., Alonso, R., Belmonte, J. A., et al. 2009, *A&A*, 494, 391
- Rouan, D., Parviainen, H., Moutou, C., et al. 2012, *A&A*, 537, A54
- Sanchis-Ojeda, R., Winn, J. N. 2011, *ApJ*, 743, 61
- Sanchis-Ojeda, R., Winn, J. N., Holman, M. J., et al. 2011, *ApJ*, 733, 127
- Scandariato, G., Maggio, A., Lanza, A. F., et al. 2013, arXiv:1301.7748
- Schwamb, M. E., Orosz, J. A., Carter, J. A. et al. 2012, submitted to *ApJ*, arXiv:1210.3612
- Seager, S., & Sasselov, D. D. 2000, *ApJ*, 537, 916
- Silva, A. V. R. 2003, *ApJ*, 585, L147
- Sing, D. K., Désert, J.-M., Lecavelier des Etangs, A., et al. 2009, *A&A*, 505, 891
- Siverd, R. J., Beatty, T. G., Pepper, J., et al. 2012, *ApJ*, 761, 123
- Shkolnik, E., Bohlender, D. A., Walker, G. A. H., Collier Cameron, A. 2008, *ApJ*, 676, 628

- Smalley, B., Anderson, D. R., Collier-Cameron, A., et al. 2012, *A&A*, 547, A61
- Solanki, S. K. 2003, *A&ARv*, 11, 153
- Southworth, J., Hinse, T. C., Jørgensen, U. G., et al., 2009, *MNRAS*, 396, 1023
- Southworth, J., Dominik, M., Jørgensen, U. G., et al., 2011, *A&A*, 527, A8
- Southworth, J., Mancini, L., Maxted, P. F. L., et al. 2012, *MNRAS*, 422, 3099
- Steffen, J. H., Fabrycky, D. C., Agol, E., et al. 2013, *MNRAS*, 428, 1077
- Strassmeier, K. G. 2009, *Astron. Astrophys. Rev.*, 17, 251
- Swain, M. R., Vasisht, G., & Tinetti, G. 2008, *Nature*, 452, 329
- Szabó, Gy. M., Szabó, R., Benkő, J. M. et al. 2011, *ApJ*, 736, L4
- Tinetti, G., et al. 2012, *Experimental Astronomy*, 34, 311
- Tregloan-Reed, J., Southworth, J., & Tappert, C. 2013, *MNRAS*, 428, 3671
- Tusnski, L. R. M., & Valio, A. 2011, *ApJ*, 743, 97
- Valenti, J. A. & Piskunov, N. 1996, *A&AS*, 118, 595
- Vogt, S. S., Hatzes A., Misch A., Kürster, M. 1999, *ApJS*, 121, 547
- Winn, J. N., Fabrycky, D., Albrecht, S., Johnson, J. A. 2010, *ApJ*, 718, L145
- Yi, S., Demarque, P., Kim, Y.-C., et al. 2001, *ApJS*, 136, 417
- Alonso, R., Brown, T. M., Torres, G., et al. 2004, *ApJ*, 613, L153
- McCullough, P. R., Stys, J. E., Valenti, J. A., et al. 2005, *PASP*, 117, 783
- Zacharias, N., Finch, C. T., Girard, T. M., et al. 2012, *VizieR Online Data Catalog*, 1322, 0Z